

Conformal decomposition in canonical general relativity

Charles H.-T. Wang^{1,2,*}

¹*School of Engineering and Physical Sciences, University of Aberdeen,
King's College, Aberdeen AB24 3UE, Scotland*

²*Space Science and Technology Department,
Rutherford Appleton Laboratory, Didcot, Oxon OX11 0QX, England*

Abstract

A new canonical transformation is found that enables the direct canonical treatment of the conformal factor in general relativity. The resulting formulation significantly simplifies the previously presented conformal geometrodynamics and provides a further theoretical basis for the conformal approach to loop quantum gravity.

PACS numbers: 04.20, 04.60

*Electronic address: c.wang@abdn.ac.uk

I. BACKGROUND

The well-known Arnowitt-Deser-Misner (ADM) canonical variables for general relativity (GR) consist of the spatial metric g_{ab} with the conjugate momentum p^{ab} . For bookkeeping purposes we will referred to these variables as the “ G -variables”. The dynamical evolution is generated by the action of the form

$$\int dt \int d^3x [p^{ab} \dot{g}_{ab} - N\mathcal{H} - N^a \mathcal{H}_a] \quad (1)$$

where the over dot denotes a t -derivative, N is the lapse function and N^a the shift vector. This action contains the the momentum (diffeomorphism) constraint:

$$\mathcal{H}_a = -2 \nabla_b p^b_a \quad (2)$$

and the Hamiltonian constraint:

$$\mathcal{H} = \mu^{-1} g_{abcd} p^{ab} p^{cd} - \mu R \quad (3)$$

where $\mu := \sqrt{\det g_{ab}}$, R is the Ricci scalar of g_{ab} and $g_{abcd} := (g_{ac}g_{bd} + g_{ad}g_{bc} - g_{ab}g_{cd})/2$. The Poisson bracket (PB) with respect to the G -variables (g_{ab}, p^{ab}) is denoted by $\{\cdot, \cdot\}^G$.

In a recent paper [1] the ADM phase space has been extended to that consisting of York’s mean curvature “time” variable $\tau := (4/3)K$, where K is the mean extrinsic curvature, with μ as momentum and conformal metric γ_{ab} with momentum π^{ab} . Based on York’s conformal decomposition of tensors, a canonical transformation has been found to relate the G -variables to the “ Γ -variables” $(\gamma_{ab}, \pi^{ab}; \tau, \mu)$ via

$$g_{ab} = \phi^4 \gamma_{ab} \quad (4)$$

$$p^{ab} = \phi^{-4} \pi^{ab} - \frac{1}{2} \phi^2 \mu_\gamma \gamma^{ab} \tau \quad (5)$$

where $\mu_\gamma := \sqrt{\det \gamma_{ab}}$ and

$$\phi := \left(\frac{\mu}{\mu_\gamma} \right)^{1/6} \quad (6)$$

is the conformal factor. Note that here ϕ is regarded as a function of the canonical variables μ and γ_{ab} . It can be seen from the above relations that a local rescaling of γ_{ab} and π^{ab} while

holding τ and μ leaves g_{ab}, p^{ab} invariant. This redundancy of the Γ -variables is offset by the “conformal constraint”:

$$\mathcal{C} = \gamma_{ab} \pi^{ab} \quad (7)$$

that generates conformal transformations through its PB with respect to the Γ -variables denoted by $\{\cdot, \cdot\}^\Gamma$. The tensor γ_{ab} will play the role of the conformal metric, whereas the original g_{ab} will stay as the physical metric.

The ADM diffeomorphism and Hamiltonian constraints then take the following forms:

$$\mathcal{H}_a = -2 \nabla_{\gamma b} \pi^b_a + \mu \tau_{,a} + 4(\ln \phi)_{,a} \mathcal{C} \quad (8)$$

$$\mathcal{H} = \frac{1}{\mu} \pi_{ab} \pi^{ab} - \frac{3}{8} \tau^2 \mu - \phi^2 \mu_\gamma R_\gamma + 8 \mu_\gamma \phi \Delta_\gamma \phi + \frac{\tau}{2} \mathcal{C} - \frac{1}{2\mu} \mathcal{C}^2. \quad (9)$$

The indices of π^{ab} and π_{ab} are raised or lowered by the conformal metric γ_{ab} and its inverse γ^{ab} respectively. Here we have used the Levi-Civita connection ∇_γ , scalar curvature R_γ and Laplacian $\Delta_\gamma := \gamma^{ab} \nabla_{\gamma a} \nabla_{\gamma b}$, associated with the conformal metric γ_{ab} .

It is useful to introduce the diffeomorphism constraint $\mathcal{C}_a$ for the Γ -variables by

$$\mathcal{C}_a := \mathcal{H}_a - 4(\ln \phi)_{,a} \mathcal{C} = -2 \nabla_{\gamma b} \pi^b_a + \mu \tau_{,a}. \quad (10)$$

Using the preservation of the PB relations by the canonical transformation from the G - to Γ -variables, the constraints $\mathcal{H}, \mathcal{C}_a$ and $\mathcal{C}$ can be explicitly shown to be first class [1, 2].

By using the relation

$$p^{ab} \dot{g}_{ab} = -\tau \dot{\mu} + \pi^{ab} \dot{\gamma}_{ab} + 4(\ln \phi) \dot{\mathcal{C}} \quad (11)$$

the canonical action for GR in the Γ -variables can be written as

$$\int dt \int d^3x [\pi^{ab} \dot{\gamma}_{ab} + \mu \dot{\tau} - N \mathcal{H} - N^a \mathcal{C}_a - \Lambda \mathcal{C}] \quad (12)$$

where Λ is a Lagrange multiplier used to effect weakly vanishing of $\mathcal{C}$. For further background information on the conformal decomposition in GR, see [1, 2, 3] and references therein.

II. CONFORMAL FACTOR AS A CANONICAL VARIABLE FOR GRAVITY

The structure of $\mathcal{H}$ in (9) is quite complicated, with ϕ being a highly nonlinear expression given in (6). If a canonical transformation can be found that retains γ_{ab} as canonical variables

while turning ϕ into a canonical variable, then the conformal decomposition for canonical GR will be greatly simplified. Below we show that this can indeed be done by explicit construction of new a set of variables satisfying these criteria. To this end, consider the quantities $\tilde{\pi}^{ab}$ and $\tilde{\pi}$ given by:

$$\tilde{\pi}^{ab} := \pi^{ab} - \frac{1}{2} \phi^6 \mu_\gamma \tau \gamma^{ab} \quad (13)$$

$$\tilde{\pi} := -6 \phi^5 \mu_\gamma \tau = -8 \phi^5 \mu_\gamma K = 4 \phi^{-1} g_{ab} p^{ab}. \quad (14)$$

From (14) we see that

$$\tau = -\frac{1}{6} \phi^{-5} \mu_\gamma^{-1} \tilde{\pi}. \quad (15)$$

Using (13) and (15) we have

$$\pi^{ab} = \tilde{\pi}^{ab} - \frac{1}{12} \phi \tilde{\pi} \gamma^{ab}. \quad (16)$$

In terms of these quantities, Eqs. (4) and (5) become simply

$$g_{ab} = \phi^4 \gamma_{ab} \quad (17)$$

$$p^{ab} = \phi^{-4} \tilde{\pi}^{ab}. \quad (18)$$

It follows that

$$-\tau \dot{\mu} = \tilde{\pi} \dot{\phi} - \phi^6 \tau \dot{\mu}_\gamma \quad (19)$$

and

$$\pi^{ab} \dot{\gamma}_{ab} = \tilde{\pi}^{ab} \dot{\gamma}_{ab} + \phi^6 \tau \dot{\mu}_\gamma. \quad (20)$$

Therefore the relation

$$\pi^{ab} \dot{\gamma}_{ab} - \tau \dot{\mu} = \tilde{\pi}^{ab} \dot{\gamma}_{ab} + \tilde{\pi} \dot{\phi} \quad (21)$$

holds and it shows that the variables $(\gamma_{ab}, \tilde{\pi}^{ab}; \phi, \tilde{\pi})$ are canonical. We shall refer to them as the “ Φ -variables” and denote the PB with respect to them by $\{\cdot, \cdot\}^\Phi$. Using (17), (18) we see the preservation of the PBs $\{\mathcal{A}, \mathcal{B}\}^G = \{\mathcal{A}, \mathcal{B}\}^\Phi$ for any $\mathcal{A}$ and $\mathcal{B}$ depending on the Φ -variables through the G -variables. In particular, the Dirac algebra for $\mathcal{H}$ and $\mathcal{H}_a$ is preserved.

From (16) we see that the conformal constraint $\mathcal{C}$ given in (7) becomes

$$\mathcal{C} = \gamma_{ab} \tilde{\pi}^{ab} - \frac{1}{4} \phi \tilde{\pi}. \quad (22)$$

The constraint $\mathcal{C}$ generates the conformal transformation in the Φ -variables via the following PB relations:

$$\{\gamma_{ab}(x), \mathcal{C}(x')\}^\Phi = \gamma_{ab}(x) \delta(x, x') \quad (23)$$

$$\{\tilde{\pi}^{ab}(x), \mathcal{C}(x')\}^\Phi = -\tilde{\pi}^{ab}(x) \delta(x, x') \quad (24)$$

$$\{\phi(x), \mathcal{C}(x')\}^\Phi = -\frac{1}{4} \phi(x) \delta(x, x') \quad (25)$$

$$\{\tilde{\pi}(x), \mathcal{C}(x')\}^\Phi = \frac{1}{4} \tilde{\pi}(x) \delta(x, x'). \quad (26)$$

From (17) and (18) we see that the physical 3-metric and its momentum are independent of on $\tilde{\pi}$ and are conformally invariant:

$$\{g_{ab}(x), \mathcal{C}(x')\}^\Phi = 0, \quad \{g_{ab}(x), \phi(x')\}^\Phi = 0 \quad (27)$$

$$\{p^{ab}(x), \mathcal{C}(x')\}^\Phi = 0, \quad \{p^{ab}(x), \phi(x')\}^\Phi = 0 \quad (28)$$

which, along with the preservation of PBs, imply

$$\{\mathcal{H}(x), \mathcal{C}(x')\}^\Phi = 0, \quad \{\mathcal{H}(x), \phi(x')\}^\Phi = 0 \quad (29)$$

$$\{\mathcal{H}_a(x), \mathcal{C}(x')\}^\Phi = 0, \quad \{\mathcal{H}_a(x), \phi(x')\}^\Phi = 0. \quad (30)$$

It follows that $\mathcal{H}, \mathcal{H}_a, \mathcal{C}$ form a set of first class constraints. To construct the explicit expressions of $\mathcal{H}$ and $\mathcal{H}_a$ in the Φ -variables, substitute (17), (18) and (22) into (2) to get

$$\mathcal{H}_a = -2 \nabla_{\gamma b} \tilde{\pi}^b_a + \tilde{\pi} \phi_{,a} + 4(\ln \phi)_{,a} \mathcal{C} \quad (31)$$

and substitute (17), (18) into (3) to get

$$\mathcal{H} = \phi^{-6} \mu_\gamma^{-1} \gamma_{abcd} \tilde{\pi}^{ab} \tilde{\pi}^{cd} - \phi^2 \mu_\gamma R_\gamma + 8 \mu_\gamma \phi \Delta_\gamma \phi \quad (32)$$

where $\gamma_{abcd} := (\gamma_{ac}\gamma_{bd} + \gamma_{ad}\gamma_{bc} - \gamma_{ab}\gamma_{cd})/2$. As with the Γ -variables, it is useful to introduce the diffeomorphism constraint in the Φ -variables as

$$\mathcal{C}_a = \mathcal{H}_a - 4(\ln \phi)_{,a} \mathcal{C} = -2 \nabla_{\gamma b} \tilde{\pi}^b_a + \tilde{\pi} \phi_{,a} \quad (33)$$

and adopt

$$\int dt \int d^3x \left[\tilde{\pi}^{ab} \dot{\gamma}_{ab} + \tilde{\pi} \dot{\phi} - N \mathcal{H} - N^a \mathcal{C}_a - \Lambda \mathcal{C} \right] \quad (34)$$

as the canonical action for GR in the Φ -variables.

It follows from the Dirac algebra for the original ADM constraints that the complete set of independent constraints $\{\mathcal{C}, \mathcal{C}_a, \mathcal{H}\}$ is indeed first class, satisfying the following algebra:

$$\{\mathcal{C}(x), \mathcal{C}(x')\}^\Phi = 0 \quad (35)$$

$$\{\mathcal{C}_a(x), \mathcal{C}_b(x')\}^\Phi = \mathcal{C}_b(x) \delta_{,a}(x, x') - (ax \leftrightarrow bx') \quad (36)$$

$$\{\mathcal{C}_a(x), \mathcal{C}(x')\}^\Phi = \mathcal{C}(x) \delta_{,a}(x, x') \quad (37)$$

$$\{\mathcal{H}(x), \mathcal{H}(x')\}^\Phi = \phi^{-4} \gamma^{ab} (\mathcal{C}_a + 4(\ln \phi)_{,a} \mathcal{C})(x) \delta_{,b}(x, x') - (x \leftrightarrow x') \quad (38)$$

$$\{\mathcal{C}(x), \mathcal{H}(x')\}^\Phi = 0 \quad (39)$$

$$\{\mathcal{C}_a(x), \mathcal{H}(x')\}^\Phi = \mathcal{H}(x) \delta_{,a}(x, x'). \quad (40)$$

The above new formulation can be applied to simplify the spin-gauge treatment in [1, 2] and to develop the conformal approach to loop quantum gravity.

Acknowledgments

I would like to thank R. Bingham, S. Carlip, J. T. Mendonça, N. O'Murchadha and J. W. York for stimulating discussions and the Aberdeen Centre for Applied Dynamics Research and the CCLRC Centre for Fundamental Physics for financial support.

-
- [1] C. H.-T. Wang, “Conformal geometrodynamics: True degrees of freedom in a truly canonical structure”, *Phys. Rev. D* **71**, 124026 (2005)
 - [2] C. H.-T. Wang, “Unambiguous spin-gauge formulation of canonical general relativity with conformorphism invariance”, *Phys. Rev. D* **72**, 087501 (2005)
 - [3] C. H.-T. Wang, “Towards conformal loop quantum gravity”, Talk given at the 4th Meeting on Constrained Dynamics and Quantum Gravity, Cala Gonone, Sardinia, Italy, 12–16 September 2005 (arXiv:gr-qc/0512023)